



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Breeding of the Rose-breasted Grosbeak at Beverly, New Jersey.—On May 2, 1896, I recorded the first arrival of this bird at Beverly, and afterwards from day to day noted a male bird whistling from the tree tops in the lawns of adjoining properties to where I live. Knowing they were not early breeders I did not pay any particular attention to him until June 1, at which time he became so noisy with his continual outbursts of song, that I concluded to investigate, and soon discovered the female building the nest. The male would accompany her about the neighborhood while gathering material, and perch on the topmost branch of a nearby tree and whistle one strain after another. After incubation commenced he became more quiet, only whistling occasionally, and going off by himself on feeding trips, as I often heard him several squares away.

The nest was situated in the topmost branches or ^{twigs} of a small maple tree, about eighteen feet above the ground, and fifteen feet from the rear of a house along the river bank within the city limits of Beverly.

As this seems very unusual, as well as the most southern record for New Jersey, so far as I can ascertain, I concluded to record the note.—J. HARRIS REED, *Beverly, N. J.*

On the Status of *Lanius robustus* Baird as a North American Bird.—Having recently had occasion to again examine the type of *Lanius robustus* Baird, which I was fortunately able to do through the kindness of the authorities of the Academy of Natural Sciences of Philadelphia, I feel more sure than ever that the bird is not North American at all. It agrees in all characters but two, namely, the larger, more strongly hooked bill and peculiar pattern of the secondaries, with *L. algeriensis*; and two examples of the latter in the National Museum collection approach it so closely in the last respect that I have little doubt it is merely an 'aberrant' specimen or possibly a local form of that species.—ROBERT RIDGWAY, *U. S. National Museum, Washington, D. C.*

***Vireo flavoviridis* in Nebraska—a Correction.**—In order to avoid confusion, it becomes necessary to state that the bird on which the note in 'The Auk' (XIII, 263), recording the capture of *Vireo flavoviridis* was founded, has been found to be *V. olivaceus*.

In explanation of how this error occurred I will state that the specimen was somewhat abnormally colored, and according to the measurements given by Ridgway (Manual, 470), had a tail at least one inch shorter than the minimum length in *V. olivaceus*. Hence it was identified as *V. flavoviridis*. Since that time I have found, however, that the measurement of the tail, 3.15–3.30, given by Ridgway is in all probability an error. All of the authorities, Coues, Baird, Brewer, and Ridgway himself in his 'Ornithology of Illinois', do not give a maximum measurement of more than 2.50.—L. BRUNER, *Lincoln, Nebraska.*